



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Notice has been sent out by the authorities that a general cleaning will be undertaken throughout all Yokohama in September under the charge of the sanitary bureau.

The sanitary condition of Yokohama is good.

Doctor Worden further reports:

Summary of dysentery, enteric fever, plague, and smallpox in Japan from January to July 10, 1909.

Dysentery.—From January of this year to July 10 there were 1,398 cases and 220 deaths. Most of the cases are reported from Aichi-ken, where there were 165 cases, with 22 deaths; 33 cases occurred in the cities and 132 cases in the rural districts. Kumamoto-ken comes next in number of cases, viz, 119 cases, with 18 deaths; 3 in the city and 116 in the rural districts. Cases were reported from all the prefectures except Saga and Miyagi. During the same period in 1908, January to July 10, 6,596 cases, with 1,565 deaths, were reported. A great decrease is thus shown over last year.

Dysentery is prevailing at present in Tokushima, Shikoku Island.

Enteric fever.—During the same period, January to July 10, of the current year 6,441 cases, with 1,466 deaths, were reported in all Japan. Tokyo had 530 cases in the city and 131 in the suburbs, with 91 deaths. In Fukuoka-ken 59 cases were reported in the cities and 808 in the rural districts, aggregating 367 in the prefecture, with 76 deaths. The smallest number, only 2 cases, with 1 death, was reported from Okinawa-ken.

Plague.—The last case of plague was reported from the province of Iyo, Shikoku Island, August 24. The governor of the prefecture has informed the home department that this case is determined as positive upon bacteriological proof. He also reports detection of some infected rats. In Formosa 1,016 cases have been reported during this year.

Smallpox.—Ninety-nine cases, with 25 deaths, were reported from all Japan. Of these 44 cases, with 12 deaths, were in Okinawa-ken. During the same period of last year there were 18,043 cases, with 5,982 deaths, reported from all Japan.

Examination of emigrants.

Number of emigrants per steamship *Nikko Maru*, for Manila, August 21: Examined 22, rejected 1, held 7.

Per steamship *Asia*, for Honolulu and San Francisco, August 28: For Honolulu, held 7; for San Francisco, held 2.

Per steamship *Kaga Maru*, September 1, for Seattle, Wash.: Rejected 2; held for observation, 4.

Report from Kobe—Inspection of vessels—Examination of emigrants—Plague—Cholera at Amagasaki.

Acting Assistant Surgeon Knight reports, August 25:

Week ended August 21. Supplemental bills of health granted to 2 steamships.

There were inspected 657 steerage passengers, and 8 steerage passengers were bathed and disinfected; their effects were disinfected